Guntersville Reservoir Watershed HUC TN06030001 TRACY CITY **Support status Fully supporting** Not supporting Not assessed Cities SOUTH PITTSBURG _o Alabama 20 Miles

Guntersville Reservoir Watershed Atlas

HUC Code: TN06030001

Counties: Franklin Grundy

Marion

Ecoregions: 68a

68b

68c

Drainage Size of Watershed: 322 square miles

Stream Miles in Watershed: 424.3
Stream Miles Fully Supporting: 133.1
Stream Miles Partially Supporting: 0.5
Stream Miles Not Supporting: 7.1
Stream Miles Not Assessed: 283.6

Lake Acres in Watershed: 1479

Lake Acres Fully Supporting: 1479 (100%)

TDEC Monitoring Stations: 20 Non-TDEC Monitoring Stations: 1


Advisories: 0

Watershed Monitoring Group: 5

Surface Water Quality in Guntersville Reservoir Watershed


Only 16 percent of the watershed is in Tennessee with the remainder in Alabama. This is a rural area with small farms and mining. Data were only available to assess 33 percent of the streams. However, 94 percent of the surveyed streams were fully supporting. Pathogens and siltation were the primary pollutant. Guntersville Reservoir is fully supporting.

This watershed has one high quality stream that is a subecoregion reference site, Crow Creek in 68c (Plateau Escarpment).



2002 Assessment of Rivers and Streams in Guntersville Reservoir Watershed

Wheeler Reservoir Watershed HUC TN06030002



Wheeler Reservoir Watershed Atlas

HUC Code: TN06030002

Counties: Franklin Giles

Lawrence Lincoln

Ecoregions: 68a

68c

71f 71g

Drainage Size of Watershed: 236 square miles

Stream Miles in Watershed: 313.3
Stream Miles Fully Supporting: 0.0
Stream Miles Partially Supporting: 24.5
Stream Miles Not Supporting: 0.0
Stream Miles Not Assessed: 288.8

Lake Acres in Watershed: None

TDEC Monitoring Stations: 19 Non-TDEC Monitoring Stations: 9


Advisories: None

Watershed Monitoring Group: 2


Surface Water Quality in Wheeler Reservoir Watershed

Eight percent of this watershed is in Tennessee. The rest is in Alabama.

Only two streams have been assessed in this watershed. Both are partially supporting. The Flint River is impaired by siltation and habitat alterations from crop production activities. This watershed is scheduled to be surveyed in Fall 2002 and 19 stations have been established. Assessment data will be included in the 2004 report.



2002 Assessment of Rivers and Streams in Wheeler Reservoir Watershed



Upper Elk River Watershed Atlas

HUC Code: TN06030003

Counties: Coffee Franklin Giles

Grundy Lincoln Marshall

Moore

Ecoregions: 68a 68c

71h 71g

Drainage Size of Watershed: 1260 square miles

Stream Miles in Watershed: 1,812.5
Stream Miles Fully Supporting: 881.4
Stream Miles Partially Supporting: 144.0
Stream Miles Not Supporting: 129.6
Stream Miles Not Assessed: 657.5

Lake Acres in Watershed: 14,504

Lake Acres Fully Supporting: 10,596 (73.1%) Lake Acres Not Supporting: 3,908 (26.9%)

TDEC Monitoring Stations: 108 Non-TDEC Monitoring Stations: 6

Advisories:


Watershed Monitoring Group: 2

Surface Water Quality in Upper Elk River Watershed (including Tims Ford and Woods Reservoirs)

Over 99 percent of the watershed is in Tennessee with a small portion in Alabama. TVA completed Tims Ford hydroelectric dam in 1970. The U.S. Air Force completed Woods Dam in 1952 to use as a source of cooling water. Both reservoirs are popular recreation areas.

Woods Reservoir is not supporting due to PCBs from contaminated sediments. Tims Ford Reservoir, as well as 76 percent of assessed streams in the watershed, is fully supporting.

Two high quality streams are subecoregion reference sites, Mud Creek in 68c (Plateau Escarpment) and Hurricane Creek in 71g (Eastern Highland Rim).



2002 Assessment of Rivers and Streams in Upper Elk River Watershed

Lower Elk River Watershed HUC TN06030004 Support status Fully supporting -PULASKI / Not supporting Partially supporting Not assessed * Cities Alabama 10 20 Miles

Lower Elk River Watershed Atlas

HUC Code: TN06030004

Counties: Giles

Lawrence Marshall

Ecoregions: 71f

71h

Drainage Size of Watershed: 718 square miles

Stream Miles in Watershed: 1,117.3
Stream Miles Fully Supporting: 524.7
Stream Miles Partially Supporting: 50.0
Stream Miles Not Supporting: 7.2
Stream Miles Not Assessed: 535.4

Lake Acres in Watershed: None

TDEC Monitoring Stations: 80 Non-TDEC Monitoring Stations: 1


Advisories: None

Watershed Monitoring Group: 2

Surface Water Quality in Lower Elk River Watershed

Seventy-six percent of the watershed is in Tennessee with the remainder in Alabama. From Tennessee, the Elk River flows into Wheeler Reservoir on the Tennessee River in Alabama.

The drainage area is primarily agricultural with row crops and pasture prevalent. Assessments in this watershed have increased from eight percent in 2000 to 52 percent for this report. Most of the newly assessed steams (90 percent) were fully supporting. Industry, municipal point sources and livestock account for the majority of impaired stream miles.



2002 Assessment of Rivers and Streams in Lower Elk River Watershed

Pickwick Reservoir Watershed HUC TN06030005 , LAWRENCEBURG PICKWICK LORETTO **Support status Fully supporting** Partially supporting Not assessed Mississippi Alabama Cities 20 40 Miles 20

Pickwick Reservoir Watershed Atlas

HUC Code: TN06030005

Counties: Hardin

Lawrence Wayne

Ecoregions: 65i

65j 71f

Drainage Size of Watershed: 639 square miles

Stream Miles in Watershed: 953.2
Stream Miles Fully Supporting: 247.1
Stream Miles Partially Supporting: 72.4
Stream Miles Not Supporting: 0.0
Stream Miles Not Assessed: 633.7

Lake Acres in Watershed: 5,840

Lake Acres Fully Supporting: 5,840 (100%)

TDEC Monitoring Stations: 81


Advisories: None

Watershed Monitoring Group: 2

Surface Water Quality in Pickwick Reservoir Watershed


Only 28 percent of the watershed is in Tennessee with the remainder in Mississippi and Alabama. Pickwick Reservoir is a TVA impoundment of the Tennessee River. Seventy-seven percent of assessed streams are fully supporting. Industry, municipal point source, and livestock are the primary pollution sources. EPA has approved organic enrichment, ammonia TMDLs on a segment of Shoal Creek (2.3 stream miles).

This watershed has four high quality streams that are subecoregion reference sites, Battle Branch in 65i (Fall Line Hills), Pompeys Branch and Dry Creek in 65j (Transition Hills), and Swanegan Branch in 71f (Western Highland Rim).



2002 Assessment of Rivers and Streams in Pickwick Reservoir Watershed

Upper Kentucky Reservoir Watershed



Upper Kentucky Reservoir Watershed Atlas

HUC Code: TN06040001

Counties: Benton Chester Decatur Hardin Humphreys Henderson

McNairy Perry Wayne

Ecoregions: 65a 65e 65i

65j 71f

Drainage Size of Watershed: 2,049 square miles

Stream Miles in Watershed: 3,435.2
Stream Miles Fully Supporting: 1,119.3
Stream Miles Partially Supporting: 153.0
Stream Miles Not Supporting: 9.8
Stream Miles Not Assessed: 2,153.1

Lake Acres in Watershed: 20,763

Lake Acres Fully Supporting: 17,500 (84.3%) Lake Acres Not Assessed: 3,263 (15.7%)

TDEC Monitoring Stations: 140 Non-TDEC Monitoring Stations: 3

Advisories: None


Watershed Monitoring Group: 3

Surface Water Quality in Upper Kentucky Reservoir Watershed


Over 98 percent of the watershed is in Tennessee with a small portion in Mississippi. Between 1963 and 1965 TVA constructed dams on the Beech River and seven tributaries for flood control and recreational use

Logging, agriculture and channelization are the primary pollution sources with siltation the most prevalent pollutant. Eighty-seven percent of assessed streams are fully supporting.

Two high quality streams are subecoregion reference sites, Right Fork Whites Creek and an unnamed tributary to Right Fork Whites Creek in 65j (Transition Hills).



2002 Assessment of Rivers and Streams in Upper Kentucky Reservoir Watershed



Upper Duck River Watershed Atlas

HUC Code: TN06040002

Counties: Bedford Coffee

Marshall Williamson

Ecoregions: 71g

71h

71i

Drainage Size of Watershed: 1553 square miles

Stream Miles in Watershed: 1,606.9
Stream Miles Fully Supporting: 668.4
Stream Miles Partially Supporting: 311.7
Stream Miles Not Supporting: 72.8
Stream Miles Not Assessed: 554.0

Lake Acres in Watershed: 3,260

Lake Acres Fully Supporting: 3,260 (100%)

TDEC Monitoring Stations: 287 Non-TDEC Monitoring Stations: 45


Advisories: 2

Watershed Monitoring Group: 3


Surface Water Quality in Upper Duck River Watershed (including Normandy Reservoir)

The entire watershed is in Tennessee. Normandy Dam, built for flood control, is TVA's largest non-power generating dam. Sixty-four percent of surveyed stream miles were fully supporting. Pathogens, nutrients, siltation and habitat alteration from agricultural activities impair the most stream miles.

Portions of the Duck River are designated as a State Scenic River. The river also provides habitat for several endangered species. Two high quality streams are subecoregion reference sites, Flat and Little Flat Creeks in 71i (Inner Nashville Basin).



2002 Assessment of Rivers and Streams in Upper Duck River Watershed



Lower Duck River Watershed Atlas

HUC Code: TN06040003

Counties: Dickson Hickman Humphreys Lawrence

Lewis Maury
Perry Wayne

Williamson

Ecoregions: 71f

71h

Drainage Size of Watershed: 736 square miles

Stream Miles in Watershed: 2,461.8
Stream Miles Fully Supporting: 1,374.0
Stream Miles Partially Supporting: 70.9
Stream Miles Not Supporting: 13.1
Stream Miles Not Assessed: 1,003.8

Lake Acres in Watershed: 13

Lake Acres Not Assessed: 13 (100%)

TDEC Monitoring Stations: 86 Non-TDEC Monitoring Stations: 24


Advisories: None

Watershed Monitoring Group: 3


Surface Water Quality in Lower Duck River Watershed

The entire watershed is in Tennessee. The area is primarily agricultural with some small towns and industry. There are also some abandoned mines. Ninety-four percent of assessed streams are fully supporting. Point source discharges (industrial and municipal), urban runoff, abandoned mines and livestock operations are sources of impairment.

This watershed has two high quality streams that are subecoregion reference sites, Wolf and Little Swan Creeks in 71f (Western Highland Rim).



2002 Assessment of Rivers and Streams in Lower Duck River Watershed



Buffalo River Watershed Atlas

HUC Code: TN06040004

Counties: Hickman Humphreys

Lawrence Lewis Perry Wayne

Ecoregions: 65j

71f

Drainage Size of Watershed: 1,823 square miles

Stream Miles in Watershed: 1,200.0
Stream Miles Fully Supporting: 616.6
Stream Miles Partially Supporting: 5.1
Stream Miles Not Supporting: 14.1
Stream Miles Not Assessed: 564.2

Lake Acres in Watershed: 349

Lake Acres Not Assessed: 349 (100%)

TDEC Monitoring Stations: 87

Advisories: None


Watershed Monitoring Group: 3

Surface Water Quality in Buffalo River Watershed

The entire watershed is in southern middle Tennessee. The Buffalo River flows into the Duck River just upstream of its confluence with the Tennessee River.

Overall water quality is good with 97 percent of assessed stream miles fully supporting designated uses.

The Tennessee General Assembly has designated portions of the Buffalo River as a State Scenic River. It is popular for canoeists and supports several commercial operators. This watershed also has one high quality stream that is a subecoregion reference site, Bush Creek in 71f (Western Highland Rim).



2002 Assessment of Rivers and Streams in Buffalo River Watershed

Lower Kentucky Reservoir Watershed **HUC TN06040005** Kentucky **PARIS** CAMDEN, WAVERLY **Support status Fully supporting** Not supporting **Partially supporting** Not assessed Cities 20 20 Miles

Lower Kentucky Reservoir Watershed Atlas

HUC Code: TN06040005

Counties: Benton Carroll Henderson Henry

Houston Humphreys

Stewart

Ecoregions: 65e 71f

74b

Drainage Size of Watershed: 1430 square miles

Stream Miles in Watershed: 2,042.6
Stream Miles Fully Supporting: 602.4
Stream Miles Partially Supporting: 186.0
Stream Miles Not Supporting: 5.9
Stream Miles Not Assessed: 1,248.3

Lake Acres in Watershed: 100,000

Lake Acres Fully Supporting: 100,000 (100%)

TDEC Monitoring Stations: 100 Non-TDEC Monitoring Stations: 2

Advisories: None


Watershed Monitoring Group: 2

Surface Water Quality in Lower Kentucky Reservoir Watershed

About 79 percent of the watershed is in Tennessee with the remainder in Kentucky. Kentucky Dam is in Kentucky.

Data were available to assess 31 percent of the stream miles; additional surveys are scheduled this fall. Seventy-six percent of assessed miles were fully supporting. Agriculture and channelization impair the most stream miles.

This watershed has one high quality stream that is a subecoregion reference site, Blunt Creek in 65e (Southeastern Plains and Hills).



2002 Assessment of Rivers and Streams in Lower Kentucky Reservoir Watershed